arXiv:1702.00807v1 [math.CO] 2 Feb 2017

ZERO-SUM INVARIANTS OF FINITE ABELIAN GROUPS

WEIDONG GAO, YUANLIN LI, JIANGTAO PENG AND GUOQING WANG

Abstract. The purpose of the article is to provide an unified way to formulate zero-sum invariants. Let G be a finite additive abelian group. Let $\mathcal{B}(G)$ denote the set consisting of all nonempty zero-sum sequences over G. For $\Omega \subset \mathcal{B}(G)$, let $d_\Omega(G)$ be the smallest integer t such that every sequence S over G of length $|S| \ge t$ has a subsequence in Ω. We provide some first results and open problems on $d_\Omega(G)$.

1. Introduction

Zero-sum theory on abelian groups can be traced back to 1960's and has been developed rapidly in recent three decades (see [2], [7] and [8]). Many invariants have been formulated in zero-sum theory and we list some of these invariants here. Let G be an additive finite abelian group. The EGZ-constant $\mathsf{s}(G)$ is the smallest integer t such that every sequence S over G of length $|S| \ge t$ has a zero-sum subsequence of length $\exp(G)$. This invariant $\mathsf{s}(G)$ comes from the so called Erdős-Ginzburg-Ziv theorem, which asserts that $\mathsf{s}(G) \le 2|G|-1$ and is regarded as one of two starting points of zero-sum theory. Another starting point of zero-sum theory involves the Davenport constant $\mathsf{D}(G)$, which is defined as the smallest integer t such that every sequence S over G of length $|S| \ge t$ has a nonempty zero-sum subsequence. This invariant $\mathsf{D}(G)$ was formulated by H. Davenport in 1965 and his motivation was to study algebraic number theory. Let $\eta(G)$ be the smallest integer t such that every sequence S over G of length $|S| \ge t$ has a zero-sum subsequence of length between 1 and $\exp(G)$, which was first introduced by Emade Boas in 1969 to study $\mathsf{D}(G)$. Let $\mathsf{E}(G)$ be the smallest integer t such that every sequence S over G of length $|S| \ge t$ has a zero-sum subsequence of length $|G|$, which was formulated by the first author in 1996. The above invariants on zero-sum sequences have been studied by many authors. To give a unified way of formulating zero-sum invariants, recently, Geroldinger, Grynkiewicz and Schmid [6] introduced the following concept. Let $\mathcal{L}$ be a finite set of positive integers, and let $\mathbf{d}_{\mathcal{L}}(G)$ be the smallest integer t such that every sequence S over G of length $|S| \ge t$ has a zero-sum subsequence of length in $\mathcal{L}$. Here we give a more unified way to formulate zero-sum invariants.

Let $\mathcal{B}(G)$ denote the set consisting of all nonempty zero-sum sequences over G. For $\Omega \subset \mathcal{B}(G)$, define $d_\Omega(G)$ to be the smallest integer t such that every sequence S over G of

2010 *Mathematics Subject Classification.* 11R27, 11B30, 11P70, 20K01.
Key words and phrases. zero-sum sequence.

length $|S| \ge t$ has a subsequence in Ω. If such t does not exist, we let $d_\Omega(G) = \infty$. Now we have

$$d_\Omega(G) = \begin{cases} \mathsf{D}(G), \text{ if } \Omega = \mathcal{B}(G); \\ \eta(G), \text{ if } \Omega = \{S \in \mathcal{B}(G) : 1 \le |S| \le \exp(G)\}; \\ \mathsf{s}(G), \text{ if } \Omega = \{S \in \mathcal{B}(G) : |S| = \exp(G)\}; \\ \mathsf{E}(G), \text{ if } \Omega = \{S \in \mathcal{B}(G) : |S| = |G|)\}; \\ \mathbf{d}_{\mathcal{L}}(G), \text{ if } \Omega = \{S \in \mathcal{B}(G) : |S| \in \mathcal{L}\}. \end{cases}$$

In this paper, we will provide some first results on $d_\Omega(G)$ and formulate some open problems. The rest of this paper is organized as follows. In Section 2, we present some necessary concepts and terminology. In Section 3 we study the way to present a given integer $t \ge \mathsf{D}(G)$ by $\Omega \subset \mathcal{B}(G)$ with $d_\Omega(G) = t$; In the final section we study the minimal $\Omega \subset \mathcal{B}(G)$ with respect to $d_\Omega(G) = t$.

2. Preliminaries

Let $\mathbb{N}$ denote the set of positive integers, $\mathbb{N}_0 = \mathbb{N} \cup \{0\}$. For a real number x, we denote by $\lfloor x \rfloor$ the largest integer that is less than or equal to x.

Throughout, all abelian groups will be written additively. By the Fundamental Theorem of Finite Abelian Groups we have

$$G \cong C_{n_1} \oplus \cdots \oplus C_{n_r},$$

where $r = \mathsf{r}(G) \in \mathbb{N}_0$ is the rank of G, $n_1, \ldots, n_r \in \mathbb{N}$ are integers with $1 < n_1 | \ldots | n_r$, moreover, $n_1, \ldots, n_r$ are uniquely determined by G, and $n_r = \exp(G)$ is the *exponent* of G. Set

$$M(G) = 1 + \sum_{i=1}^{r} (n_i - 1).$$

Let $G_0 \subset G$. For $g_1, \ldots, g_l \in G_0$(repetition allowed), we call $S = g_1 \cdot \ldots \cdot g_l$ a *sequence* over G_0. We write sequences S in the form

$$S = \prod_{g \in G} g^{\mathsf{v}_g(S)} \text{ with } \mathsf{v}_g(S) \in \mathbb{N}_0 \text{ for all } g \in G_0.$$

We call $\mathsf{v}_g(S)$ the *multiplicity* of g in S.

For $S = g_1 \cdot \ldots \cdot g_l = \prod_{g \in G_0} g^{\mathsf{v}_g(S)}$, we call

- $|S| = l = \sum_{g \in G_0} \mathsf{v}_g(S) \in \mathbb{N}_0$ the *length* of S.
- $\sigma(S) = \sum_{i=1}^{l} g_i = \sum_{g \in G_0} \mathsf{v}_g(S) g \in G_0$ the *sum* of S.
- S is a *zero-sum sequence* if $\sigma(S) = 0$.
- S is a *short zero-sum sequence* if it is a zero-sum sequence of length $|S| \in [1, \exp(G)]$

We denote by $\mathcal{B}(G_0)$ the set of all nonempty zero-sum sequences over G_0, by $\mathcal{A}(G_0)$ the set of all minimal zero-sum sequences over G_0.

We say two sequences S and T over G have the same form if and only if $\mathsf{v}_g(S) = \mathsf{v}_g(T)$ for every $g \in G$.

3. Representing the same invariant t by different Ω with $d_\Omega(G) = t$

We first state some basic properties on $d_\Omega(G)$.

Proposition 3.1. *Let $\Omega, \Omega' \subset \mathcal{B}(G)$.*

1. *$d_\Omega(G) < \infty$ if and only if for every element $g \in G$, $g^{k\,\mathrm{ord}(g)} \in \Omega$ for some positive integer $k = k(g)$*
2. *If $\Omega \subset \Omega'$ then $d_{\Omega'}(G) \le d_\Omega(G)$.*
3. *If S_1 is a proper subsequence of S and both S and S_1 belong to Ω, then $d_{\Omega\setminus\{S_1\}}(G) = d_\Omega(G)$.*
4. *For every positive integer $t \in [\mathsf{D}(G), \infty)$, there is an $\Omega \subset \mathcal{B}(G)$ such that $t = d_\Omega(G)$.*

Proof. 1. Necessity. Let $t = d_\Omega(G) < \infty$. For every $g \in G$, the sequence g^t has a nonempty zero-sum subsequence S in Ω. Since S is zero-sum, we infer that it has the form $g^{k\,\mathrm{ord}(g)}$ for some positive integer k. This proves the necessity.

Sufficiency. If for every $g \in G$ there is a positive integer k_g such that $g^{k_g\,\mathrm{ord}(g)} \in \Omega$, then clearly $d_\Omega(G) \le 1 + \sum_{g\in G}(k_g\,\mathrm{ord}(g) - 1)$ completing the proof of Conclusion 1.

2. The result holds obviously.

3. The result follows from 2.

4. Let $G^* = G \setminus \{0\}$. For each positive integer $t \ge \mathsf{D}(G)$. Let $\Omega = \{0^{t-\mathsf{D}(G)+1}\} \cup \mathcal{B}(G^*)$. It is easy to see that $t = d_\Omega(G)$, completing the proof. □

Although Conclusion 4 of the above proposition asserts that for every positive $t \ge \mathsf{D}(G)$, there is an $\Omega \subset \mathcal{B}(G)$ such that $t = d_\Omega(G)$, it does not give us much information on the invariant t. For some classical invariants t, we hope to find some special $\Omega \in \mathcal{B}(G)$ with $d_\Omega(G) = t$ to help us understand t better. We say a sequence S over G a *weak-regular* sequence if $\mathsf{v}_g(S) \le \mathrm{ord}(g)$ for every $g \in G$. We say $\Omega \subset \mathcal{B}(G)$ is *weak-regular* if very sequence $S \in \Omega$ is weak-regular. Define $Vol(G)$ to be the set of all positive integer $t \in [\mathsf{D}(G), 1 + \sum_{g\in G}(\mathrm{ord}(g) - 1)]$ such that $t = d_\Omega(G)$ for some weak-regular $\Omega \subset \mathcal{B}(G)$.

Remark 3.2. A sequence over G is called a regular sequence if $|S_H| \le |H| - 1$ for every proper subgroup H of G, where S_H denote the subsequence consisting of all terms of S in H. The concept of regular sequences was introduced by Gao, Han and Zhang [3] quite recently.

Question 1. Does $Vol(G) = [\mathsf{D}(G), 1 + \sum_{g\in G}(\mathrm{ord}(g) - 1)]$ hold for any finite abelian group G?

Let $\Omega \subset \mathcal{B}(G)$, we say a sequence S over G is Ω-*free* if S has no subsequence in Ω.

Lemma 3.3. *Let $\Omega \in \mathcal{B}(G)$ with $d_\Omega(G) = t \in [\mathsf{D}(G), 1 + \sum_{g\in G}(\mathrm{ord}(g) - 1)]$. If there is an Ω-free sequence T over G of length $|T| = t - 1$ such that $\mathsf{v}_g(T) \le \mathrm{ord}(g) - 1$ for every $g \in G$, then $d_\Omega(G) = t \in Vol(G)$.*

Proof. If every sequence in Ω is weak regular, then by the definition of $Vol(G)$ we know that $t \in Vol(G)$. Otherwise, take an $S \in \Omega$ such that $\mathsf{v}_g(S) \geq \mathrm{ord}(g) + 1$ for some $g \in G$. Let $\Omega_1 = (\Omega \setminus \{S\}) \cup \{g^{\mathrm{ord}(g)}\}$. It is easy to see that $d_{\Omega_1}(G) \leq d_\Omega(G) = t$. On the other hand, T is also Ω_1-free. Therefore, $d_{\Omega_1}(G) \geq t$. Hence, $d_{\Omega_1}(G) = t$. Continuing the same process above we finally get an Ω_k which is weak regular such that $d_{\Omega_k}(G) = t \in Vol(G)$. □

Proposition 3.4. *For every finite abelian group G we have*

$$\mathsf{D}(G) \in Vol(G), \mathsf{D}(G) + 1 \in Vol(G) \text{ and } \eta(G) \in Vol(G).$$

Proof. Let $\Omega = \mathcal{B}(G)$. Then, $d_\Omega(G) = \mathsf{D}(G)$. Let T be a zero-sum-free sequence (or equivalently, Ω-free) over G of length $|T| = \mathsf{D}(G) - 1 = d_\Omega(G) - 1$. Clearly, T is weak regular and $\mathsf{D}(G) = d_\Omega(G) \in Vol(G)$ by Lemma 3.3.

To prove $\mathsf{D}(G) + 1 \in Vol(G)$, let $\Omega = \{S \in \mathcal{B}(G), |S| < \mathsf{D}(G)\} \cup \{g^{\mathrm{ord}(G)}, g \in G\}$. It is easy to see that $d_\Omega(G) = \mathsf{D}(G) + 1$. Let $\Omega' = \{S \in \mathcal{A}(G), |S| < \mathsf{D}(G)\} \cup \{g^{\mathrm{ord}(G)}, g \in G\}$. Clearly, $d_{\Omega'}(G) = d_\Omega(G) = \mathsf{D}(G) + 1$ and $\mathsf{D}(G) + 1 \in Vol(G)$ follows from Ω' is weak-regular.

To prove $\eta(G) \in Vol(G)$, let $\Omega = \{S \in \mathcal{B}(G), |S| \leq \exp(G)\}$. It is easy to see that $d_\Omega(G) = \eta(G)$. Let T be any Ω-free sequence of the maximal length $|T| = \eta(G) - 1$. Clearly, $\mathsf{v}_g(T) \leq \mathrm{ord}(g) - 1$ for every $g \in G$ and hence $\eta(G) \in Vol(G)$ by Lemma 3.3 again. □

4. The minimal Ω with respect to $d_\Omega(G) = t$ for some given t

For a given $t \geq \mathsf{D}(G)$ and an $\Omega \subset \mathcal{B}(G)$ with $d_\Omega(G) = t$. We say that Ω is minimal respect to $d_\Omega(G) = t$ if $d_{\Omega'}(G) > t$ for every proper subset $\Omega' \subsetneq \Omega$.

By Proposition 3.1 (3) we have the following.

Proposition 4.1. *Let $t \geq \mathsf{D}(G)$ be an integer, and let $\Omega \subset \mathcal{B}(G)$ be minimal with respect to $d_\Omega(G) = t$. Then, for every $B, B' \in \Omega$ with $B \neq B'$ we have $B' \nsubseteq B$.*

Question 2. Let $\Omega \in \mathcal{B}(G)$ be minimal with respect to $d_\Omega(G) = \mathsf{D}(G)$. What can be said about Ω?

For Question 2, $d_\Omega(G) = \mathsf{D}(G)$ if $\Omega = \mathcal{A}(G)$, but we can not say that $\mathcal{A}(G)$ is smallest. Let p be a prime. Let $G = C_p$. Let Ω_1 be the set consisting of all minimal zero-sum sequences over G with index 1. If the Lemke-Kleitman's conjecture is true for p, i.e., every sequence over G of length p has a subsequence with index 1, then Ω_1 is a proper subset of $\mathcal{A}(G)$ with $d_{\Omega_1}(G) = \mathsf{D}(G) = p$. So if $n = p \geq 7$ is a prime and the Lemke-Kleitman's conjecture mentioned above is true for p, then $\Omega = \mathcal{A}(G)$ is not minimal with respect to $d_\Omega(G) = \mathsf{D}(G) = p$. In fact, except for some small primes p, we don't know any minimal $\Omega \subset \mathcal{B}(G)$ with respect to $d_\Omega(G) = \mathsf{D}(G) = p$. For sufficiently large t, we can find minimal Ω with respect to $d_\Omega(G) = t$.

Proposition 4.2. *Let G be a finite abelian group and $t \geq 1 + \sum_{g\in G}(\mathrm{ord}(g) - 1)$ be a integer, and let $\Omega = \{g^{\mathrm{ord}(g)}, g \in G\} \cup \{0^{t-\sum_{g\in G}(\mathrm{ord}(g)-1)}\}$. Then Ω is minimal with respect to $d_\Omega(G) = t$.*

Proof. Clearly.

□

For a given $t \in [\mathsf{D}(G), \sum_{g\in G}(\mathrm{ord}(g) - 1)]$ not too close to $\sum_{g\in G}(\mathrm{ord}(g) - 1)$, it could be very difficult to find a minimal $\Omega \subset \mathcal{B}(G)$ with respect to $d_\Omega(G) = t$. We provide one more example here:

Proposition 4.3. *Let $n \geq 5$ be an odd integer. Then $\Omega = \{T \in \mathcal{B}(C_n)|\ |T| = n\}$ is not minimal with respect to $d_\Omega(G) = 2n - 1$.*

Proof. Let $S_0 = \prod_{i=0}^{n-1} i$. Since n is odd, we have $S_0 \in \Omega$. Let $\Omega_1 = \Omega \setminus \{S_0\}$. Clearly, $d_{\Omega_1}(G) \geq d_\Omega(G) = 2n - 1$. So, it suffices to prove $d_{\Omega_1}(G) \leq 2n - 1$. Let S be an arbitrary sequence over G with length $|S| = 2n - 1$. We need to show S contains a zero-sum subsequence in Ω_1. Since $d_\Omega(G) = 2n-1$, we may assume that all zero-sum subsequences of S with length n is of the same form as $S_0 = \prod_{i=0}^{n-1} i$. It follows that j occurs exactly once in S for some $j \in [0, n-1]$. Consider the sequence $S_1 = S \setminus \{j\}$. Then, $|S_1| = 2n-2$ and S_1 contains $n-1 > 2$ distinct elements. By recalling the well known fact that a sequence over C_n with length $2n-2$ having no zero-sum subsequence of length n if and only if the sequence consists of two distinct elements with each appearing $n-1$ times, we derive that S_1 and hence S has a zero-sum subsequence in Ω_1. □

We say a zero-sum sequence S is *essential* with respect to some $t \geq \mathsf{D}(G)$ if every $\Omega \in B(G)$ with $d_\Omega(G) = t$ contains S. For examples, for $G = C_n$ and $t = n$, the zero-sum sequence $g(-g)$ is essential with respect to n for every generator g of C_n. For any finite abelian group G, every minimal zero-sum sequence over G of length $\mathsf{D}(G)$ is essential with respect to $\mathsf{D}(G)$.

A natural research problem is to determine the smallest integer t such that there is no essential zero-sum sequence with respect to t, denote it by $q(G)$.

Let $q'(G)$ be the smallest integer t such that, every sequence S over G of length $|S| \geq t$ has two nonempty zero-sum subsequences with different forms.

Question 3. Does $q(G) = q'(G)$ for any finite abelian group G?

Relating to $q'(G)$ we present two invariants below.

Let $disc(G)$ denote the smallest integer t such that every sequence S over G of length $|S| \geq t$ has two nonempty zero-sum subsequences with different lengths. The invariants $disc(G)$ was formulated by Girard [9] and has been studied recently in [4] and [5].

Let $\mathsf{D}_2(G)$ denote the smallest integer t such that every sequence over G has two disjoint nonempty zero-sum subsequences. Clearly,

$$q'(G) \leq disc(G) \leq \mathsf{D}_2(G)$$

for every finite abelian group G.

$\mathsf{D}_2(G)$ was first introduced by H. Halter-Koch in [10] and was studied recently by Plagne and Schmid in [11].

We close this section with the following theorem.

Theorem 4.4. *If G is a finite abelian group then $q(G) \le \mathsf{D}_2(G)$.*

Proof. Let $G^* = G \setminus \{0\}$. Let $k \ge \mathsf{D}_2(G)$ be a integer and we only need to show that there exist two disjoint subsets $\Omega, \Omega' \subset \mathcal{B}(G)$ such that $d_\Omega(G) = k = d_{\Omega'}(G)$. Let $\Omega = \{0^{k-\mathsf{D}(G)+1}\} \cup \mathcal{A}(G^*)$ and let $\Omega' = \{0^{k-\mathsf{D}_2(G)+1}\} \cup \{\mathcal{B}(G^*) \setminus \mathcal{A}(G^*)\}$.

Note that a minimal zero-sum sequence over G of length $\mathsf{D}(G)$ has no two disjoint nonempty zero-sum subsequences, so we obtain that $\mathsf{D}_2(G) > \mathsf{D}(G)$. Thus,

$$\Omega \cap \Omega' = \emptyset.$$

On the other hand, it is easy to see that $d_\Omega(G) = k = d_{\Omega'}(G)$ and we are done.

□

Acknowledgements. This work was supported by the PCSIRT Project of the Ministry of Science and Technology, the National Science Foundation of China, and a Discovery Grant from the Natural Science and Engineering Research Council of Canada.

References

[1] P. Erdős, G. Ginzburg and A. Ziv, *Theorem in the additive number theory*, Bull. Res. Council Israel, 10F(1961) 41-43.

[2] W. Gao and A. Geroldinger, *Zero-sum problems in finite abelian groups: a survey*, Expo. Math. **24** (2006), 337-369.

[3] W.D. Gao, D.C. Han and H.B. Zhang, *On additive Bases II*, Acta Arith. 168 (2015), 247-267.

[4] W.D.Gao, P.P. Zhao and J.J. Zhuang, *Zero-sum subsequences of distinct lengths*, Int. J. Number Theory, 11 (2015).

[5] W.D. Gao, Y.L. Li, P.P. Zhao and J.J. Zhuang, *ON sequences over a finite abelian group with zero-sum Ssubsequences of forbidden lengths*, Colloq. Math., 144(2016) 21-44.

[6] A. Geroldinger, D. Grynkiewicz and W. Schmid, *Zero-sum problems with congruence conditions*, Acta Math. Hungar., 131(2011) 323-345.

[7] A. Geroldinger, *Additive group theory and non-unique factorizations*, Combinatorial Number Theory and Additive Group Theory (A. Geroldinger and I. Ruzsa, eds.), Advanced Courses in Mathematics CRM Barcelona, Birkhäuser, 2009, pp. 1 – 86.

[8] A. Geroldinger and F. Halter-Koch, *Non-Unique Factorizations. Algebraic, Combinatorial and Analytic Theory*, Pure and Applied Mathematics, vol. 278, Chapman & Hall/CRC, 2006.

[9] B. Girard, *On the existence of zero-sum subsequences of distinct lengths.* Rocky Mountain J. Math. 42(2012) 583-596.

[10] F. Halter-Koch, *A generalization of Davenport's constant and its arithmetical applications.* Colloq. Math. 63(1992) 203-210.

[11] A. Plagne and W.A. Schmid, *An application of coding theory to estimating Davenport constants.* Des. Codes Cryptogr. 61 (2011) 105-118.

Center for Combinatorics and LPMC-TJKLC, Nankai University, Tianjin, 300071, P.R. China

E-mail address: wdgao@nankai.edu.cn,

Department of Mathematics, Brock University, St. Catharines, Ontario, Canada L2S 3A1
E-mail address: yli@brocku.ca

College of Science, Civil Aviation University of China, Tianjin, 300300, P.R. China
E-mail address: jtpeng1982@aliyun.com

Department of Mathematics, Tianjin Polytechnic University, Tianjin, 300387, P. R. China
E-mail address: gqwang1979@aliyun.com